A Larger Equational Proof on Lists

A Law of Reverse

For a more difficult example, let's consider the `reverse` function.

We pick its inefficient definition, because its more amenable to equational proofs:

```
      Nil.reverse  =  Nil                       // 1st clause
(x :: xs).reverse  =  xs.reverse ++ List(x)     // 2nd clause
```

We'd like to prove the following proposition

```
xs.reverse.reverse  =  xs
```

Proof

By induction on `xs`. The base case is easy:

```
Nil.reverse.reverse
=   Nil.reverse             // by 1st clause of reverse
=   Nil                     // by 1st clause of reverse
```

Proof

By induction on `xs`. The base case is easy:

```
Nil.reverse.reverse
=   Nil.reverse             // by 1st clause of reverse
=   Nil                     // by 1st clause of reverse
```

For the induction step, let's try:

```
(x :: xs).reverse.reverse
=   (xs.reverse ++ List(x)).reverse // by 2nd clause of reverse
```

Proof

By induction on `xs`. The base case is easy:

```
Nil.reverse.reverse
=   Nil.reverse             // by 1st clause of reverse
=   Nil                     // by 1st clause of reverse
```

For the induction step, let's try:

```
(x :: xs).reverse.reverse
=   (xs.reverse ++ List(x)).reverse // by 2nd clause of reverse
```

We can't do anything more with this expression, therefore we turn to the right-hand side:

```
x :: xs
=   x :: xs.reverse.reverse         // by induction hypothesis
```

Both sides are simplified in different expressions.

To Do

We still need to show:

```
(xs.reverse ++ List(x)).reverse  =  x :: xs.reverse.reverse
```

Trying to prove it directly by induction doesn't work.

We must instead try to *generalize* the equation. For *any* list `ys`,

```
(ys ++ List(x)).reverse  =  x :: ys.reverse
```

This equation can be proved by a second induction argument on `ys`.

Auxiliary Equation, Base Case

```
(Nil ++ List(x)).reverse       // to show: =  x :: Nil.reverse
```

Auxiliary Equation, Base Case

```
(Nil ++ List(x)).reverse       // to show: =  x :: Nil.reverse

=   List(x).reverse            // by 1st clause of ++
```

Auxiliary Equation, Base Case

```
(Nil ++ List(x)).reverse       // to show: =  x :: Nil.reverse

=   List(x).reverse            // by 1st clause of ++

=   (x :: Nil).reverse         // by definition of List
```

Auxiliary Equation, Base Case

```
(Nil ++ List(x)).reverse       // to show: =  x :: Nil.reverse

=   List(x).reverse            // by 1st clause of ++

=   (x :: Nil).reverse         // by definition of List

=   Nil ++ (x :: Nil)          // by 2nd clause of reverse
```

Auxiliary Equation, Base Case

```
(Nil ++ List(x)).reverse       // to show: =  x :: Nil.reverse

=   List(x).reverse            // by 1st clause of ++

=   (x :: Nil).reverse         // by definition of List

=   Nil ++ (x :: Nil)          // by 2nd clause of reverse

=   x :: Nil                   // by 1st clause of ++
```

Auxiliary Equation, Base Case

```
(Nil ++ List(x)).reverse      // to show: =  x :: Nil.reverse

=   List(x).reverse           // by 1st clause of ++

=   (x :: Nil).reverse        // by definition of List

=   Nil ++ (x :: Nil)         // by 2nd clause of reverse

=   x :: Nil                  // by 1st clause of ++

=   x :: Nil.reverse          // by 1st clause of reverse
```

Auxiliary Equation, Inductive Step

```
((y :: ys) ++ List(x)).reverse          // to show: =  x :: (y :: ys).reverse
```

Auxiliary Equation, Inductive Step

```
((y :: ys) ++ List(x)).reverse          // to show: =  x :: (y :: ys).reverse

=   (y :: (ys ++ List(x))).reverse      // by 2nd clause of ++
```

Auxiliary Equation, Inductive Step

```
((y :: ys) ++ List(x)).reverse             // to show: =  x :: (y :: ys).reverse

=   (y :: (ys ++ List(x))).reverse         // by 2nd clause of ++

=   (ys ++ List(x)).reverse ++ List(y)     // by 2nd clause of reverse
```

Auxiliary Equation, Inductive Step

```
((y :: ys) ++ List(x)).reverse           // to show: =  x :: (y :: ys).reverse

=   (y :: (ys ++ List(x))).reverse       // by 2nd clause of ++

=   (ys ++ List(x)).reverse ++ List(y)   // by 2nd clause of reverse

=   (x :: ys.reverse) ++ List(y)         // by the induction hypothesis
```

Auxiliary Equation, Inductive Step

```
((y :: ys) ++ List(x)).reverse             // to show: =  x :: (y :: ys).reverse

=   (y :: (ys ++ List(x))).reverse         // by 2nd clause of ++

=   (ys ++ List(x)).reverse ++ List(y)     // by 2nd clause of reverse

=   (x :: ys.reverse) ++ List(y)           // by the induction hypothesis

=   x :: (ys.reverse ++ List(y))           // by 1st clause of ++
```

Auxiliary Equation, Inductive Step

```
  ((y :: ys) ++ List(x)).reverse          // to show: =  x :: (y :: ys).reverse

=    (y :: (ys ++ List(x))).reverse       // by 2nd clause of ++

=    (ys ++ List(x)).reverse ++ List(y)   // by 2nd clause of reverse

=    (x :: ys.reverse) ++ List(y)         // by the induction hypothesis

=    x :: (ys.reverse ++ List(y))         // by 1st clause of ++

=    x :: (y :: ys).reverse               // by 2nd clause of reverse
```

This establishes the auxiliary equation, and with it the main proposition.

Exercise (Open-Ended, Harder)

Prove the following distribution law for `map` over concatenation.

For any lists `xs`, `ys`, function `f`:

```
(xs ++ ys) map f  =  (xs map f) ++ (ys map f)
```

You will need the clauses of $++$ as well as the following clauses for `map`:

```
      Nil map f  =  Nil
(x :: xs) map f  =  f(x) :: (xs map f)
```